

Northwest Health Care Associates

JUVÉDERM™ Consent

Indications

JUVÉDERM™ Ultra injectable gel are injected into areas of facial tissue where moderate to severe facial wrinkles and folds occur. It temporarily adds volume to the skin and subcutaneous tissues may give the appearance of a smoother skin surface and may help smooth moderate to severe facial wrinkles and folds. Correction is temporary; therefore, touch-up injections as well as repeat injections are usually needed to maintain optimal correction. Less material (about half the amount) is usually needed for repeat injections. Most patients need one or possibly two treatments to achieve optimal wrinkle smoothing. The results may last as long as 9 months to 1 year. **Initial**_____

Side Effects and Complications

Most side effects are mild or moderate in nature, and their duration is short lasting (7 days or less). The most common side effects include, but are not limited to, temporary injection-site reactions such as : redness, pain/tenderness, firmness, swelling, lumps/bumps, bruising, itching, infection and discoloration. In the first 24 hours after injection, you should avoid strenuous exercise, extensive sun or heat exposure, and alcoholic beverages. Exposure to any of the above may cause temporary redness, swelling, and/or itching at the injection sites. If there is swelling, you may need to place an ice pack over the swollen area. You should ask your physician when makeup may be applied after your treatment. Be sure to report any redness and/or visible swelling that lasts for more than a few days, or any other symptoms that cause you concern. **Initial**_____

Contraindications

JUVÉDERM™ Ultra injectable gel should not be used if you have:

- Severe allergies marked by a history of anaphylaxis or history or presence of multiple severe allergies
- A history of allergies to Gram-positive bacterial proteins **Initial**_____

The following are important treatment considerations for you to discuss with us and understand in order to help avoid unsatisfactory results and complications:

- **Please inform us prior to treatment:** If you are using substances that can prolong bleeding, such as aspirin or ibuprofen, as with any injection, may experience increased bruising or bleeding at the injection site.
- **Please inform us prior to treatment:** If you are on immunosuppressive or therapy used to decrease the body's immune response, as there may be an increased risk of infection
- **Please inform us prior to treatment:** If you are pregnant or breastfeeding,
- **Please inform us prior to treatment:** If you have history of excessive scarring (eg, hypertrophic scarring and keloid formations) and pigmentation disorders. **Initial**_____

If laser treatment, chemical peeling, or any other procedure based on active dermal response is considered after treatment with JUVÉDERM™ Ultra injectable gel, there is a possible risk of an inflammatory reaction at the treatment site. **Initial**_____

The safety and effectiveness of JUVÉDERM™ Ultra injectable gel for the treatment of areas other than facial wrinkles and folds (such as lips) have not been established in controlled clinical studies. **PATIENT'S ACCEPTANCE OF RISKS Initial**_____

I have read the above information and have discussed it with my physician. I understand that it is impossible to be informed of every possible complication that may occur. No guarantees about results have been made. By signing below, I agree that my doctor has answered all of my questions and that I understand and accept the risks, benefits, and alternatives of JUVÉDERM™ Ultra **Initial**_____

Please Print Name

Date

Patient Signature

Date